

Emerging Infectious Disease Surveillance Tool (SRI/MERS/EBOLA)



Limited Use License

By using these materials you (the “User”) agree to the following terms and conditions. The User is hereby granted a non-transferable, non-exclusive, revocable, perpetual, limited use license in the following materials. The User shall not, in whole or in part, alter or change the materials. The User acknowledges and understands that the International Academies of Emergency Dispatch is the sole and exclusive owner of the copyrights and other intellectual property associated with the materials and all derivatives therefrom. The User agrees to keep any of these materials current as regularly issued by the IAED. These materials are protected by United States and International copyright laws and treaties. The User shall not use these materials commercially or for any monetary gain. They cannot be incorporated into any 3rd party products, computers or CADs, modified in any way, or redistributed for any other uses. In no event shall the International Academies of Emergency Dispatch be liable for damages of any kind associated with the use of these materials.

This Protocol developed and approved by the IAED's CBRN Fast Track Committee of the Council of Standards.

Listen carefully and tell me if s/he has any of the following symptoms:

- difficulty breathing or shortness of breath
- persistent cough
- measured body temperature > 101.5° F (38.6° C)
- chills
- unusual sweats
- hot to the touch in room temperature
- unusual total body aches
- headache
- sore throat
- nasal congestion (blocked nose)
- runny or stuffy nose
- recent onset of any diarrhea, vomiting, or bloody discharge from the mouth or nose
- abdominal or stomach pain
- unusual (spontaneous/non-traumatic) bleeding from any area of the body
- contact with someone with the flu or flu-like symptoms (if so, when?)

continued on reverse side

Ask only in early phases when new flu, respiratory illness, or hemorrhagic fever is emerging from specific areas:

- traveled in the last 21 days (if so, where?) Note: (If travel timeframe questionable) Was it roughly within the past month?

Ask only if a higher-risk exposure is suspected (close contact with sick persons, dead bodies, or exotic African animals):

- needlestick, scalpel cut, or similar injury in treating or caring for Ebola patients
- blood or body fluid exposure to eyes, nose, or mouth (mucous membranes) in treating or caring for Ebola patients
- skin contact with, or exposure to, blood or body fluids of an Ebola patient
- direct contact with a dead body without use of personal protective equipment in a country where an Ebola outbreak is occurring
- handling of bats, rodents, or non-human primates in or recently received from Africa

Infection Prevention Instructions:

- (Keep isolated)** From now on, **don't allow** anyone to come in **close contact** with her/him.